

**LEAFLET FOR:**

**CALCIUM CHLORIDE solution for injection 50 mL**  
**CALCIUM CHLORIDE solutio pro injectionibus 50 mL**  
**For big and small ruminant animals, horses, swine and dogs.**

**1. NAME AND ADDRESS OF THE AUTHORIZATION HOLDER AND OF THE PRODUCER (IF THEY DIFFERENT)**

Marketing authorization holder and producer:

VETPROM AD, 2400, Radomir, Bulgaria, 26, Otets Paisiy Str.

**2. NAME OF THE VETERINARY MEDICINAL PRODUCT**

CALCIUM CHLORIDE solution for injection 50 mL

CALCIUM CHLORIDE solutio pro injectionibus 50 mL

**3. ACTIVE SUBSTANCE AND EXCIPIENTS**

**Active Substance:**

3.37 g Calcium chloride dihydrate is contained in 50 mL of the medicinal product

**Excipient:**

Water for injections (solvent).

**4. THERAPEUTIC INDICATIONS**

In birth paresis – before and after giving birth, in tetany of various origins – milk, transport, barn or grass; in eclampsia in swine and dogs; in allergic conditions – serum sickness, urticarial, aseptic total pododermatitis; in toxicosis, afterbirth hemoglobinuria, hemorrhages, petechial fever, hemorrhagic diathesis, carbon tetrachloride intoxication; for avoiding the excitatory stage when awakening from narcosis.

**5. CONTRAINDICATIONS**

The product is not used in thrombosis predisposition and in hypercalcemia.

**6. ADVERSE REACTIONS**

In case of administration outside the vane tissue necrosis develops - this can be prevented by 10 % sodium sulfate solution injected in the tissues.

**7. ANIMAL SPECIES, FOR WHICH THIS VMP IS DESIGNED**

For big and small ruminant animals, horses, swine and dogs.

## **8. RECOMMENDED DOSES FOR EACH OF THE ANIMAL SPECIES, ADMINISTRATION ROUTE**

**Administration:** The administration is strictly limited to slow intravenous injection!

**Dose:** The medicinal product is administered 1 time daily for several days:

- For big ruminant animals and horses – 150 to 200 mL;
- For small ruminant animals and sheep – 30 to 50 mL;
- For dogs - 0.1 to 1 mL/kg b.w.

## **9. ADVICE FOR PROPER ADMINISTRATION**

Always use CALCIUM CHLORIDE solution for injection 50 mL exactly as you are old by your veterinarian. If in doubt, ask your veterinarian.

## **10. WITHDRAWAL PERIOD**

Zero /0/ days.

## **11. SPECIAL PRECAUTIONS FOR STORAGE OF THE VMP**

Do not store above 25°C.

Protect from light.

Keep in dry place.

Keep the medication out of sight and reach of children!

## **12. SPECIAL PRECAUTIONS FOR STORAGE OF THE VMP**

Calcium can penetrate the placental barrier and can be excreted with milk.

## **13. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED VETERINARY MEDICINAL PRODUCT OR WASTE MATERIALS DERIVED FROM THE USE OF SUCH PRODUCTS**

The VMP should be destroyed in accordance with requirement of the local laws.

## **14. DATE OF THE LAST UPDATE OF THE TEXT**

SEPTEMBER 2010

## **15. ADDITIONAL INFORMATION**

Please, contact the local representative of the authorization holder for any information concerning this veterinary medicinal product.

VETPROM AD,

2400, Radomir, Bulgaria, 26, Otets Paisiy Str.

Tel.: 0777/8-24-93, 8-00-19; Fax. 0777/8-23-91

E-mail: [vetprom@abv.bg](mailto:vetprom@abv.bg)