

LEAFLET

ICHTAMMOL 10% ointment

ICHTHAMMOL VETPROM 10% ointment

Cattle, horses, donkeys, swines, sheep, goats and dogs

1. NAME AND CONSTANT ADDRESS OF THE LICENSEE FOR USE AND THE PRODUSER IF THEY ARE DIFFERENT

VETPROM AG, 2400 Radomir, Bulgaria, ul.“Otec Paisii” № 26

tel.: 0777/8-24-93, 8-02-68; fax: 0777/8-23-91

E-mail: vetprom@abv.bg

2. NAME OF THE VETERINARY AND MEDICAL PRODUCT

ICHTHAMMOL VETPROM 10% ointment

3. CONTENT OF THE ACTIVE SUBSTANCE AND THE EXCIPIENTS

50 g contain:

Active Substance: Ichthammol 5.0 g

Excipients: Adeps Suillus, White Soft Paraffin

4. THERAPEUTIC INDICATIONS

In case of superficial and deeper suppurative inflammable illnesses of the skin and subcutaneous connecting tissue- furuncles , carbuncles, flegmones , abscesses, ulcers, fistulas, suppurative wounds, eczemas, suppurative dermatitis, burnings, frozen states . In illnesses of bones and joints, like suppurative bursitis and tendovaginitis, osteomyelitis after osteosynthesis and other.

5. CONTRAINDICATIONS

Not to be used in case of hypersensitivity to the active substance or to any of the excipients.

Not to be used in animals of family Cats.

6. UNFAVOURABLE REACTIONS The Appearance of skin rashes is possible. The frequency of the unfavourable reactions is defined according to the next classification:

- very frequent (more than 1 to 10 animals, appearing unfavourable reactions through the course of one treatment)
- frequent (more than 1, but less than 10 animals from 100 ones)
- not frequent (more than one , but less than 10 animals from 1,000 ones)

- rare (more than 1, but less than 10 from 10,000 animals)

- very rare (less than 1 animal from 10,000 animals, including isolated cases).

If you notice any serious influence or other effects, as a result of the use of this VMP, that are undescribed in this leaflet, please suddenly inform your veterinary doctor.

7. KINDS OF ANIMALS, TO WHICH THIS VMP IS DESIGNED

Cattle, horses, donkeys, swines, sheep, goats and dogs.

8. DOSAGE FOR EVERY KIND OF ANIMAL, METHOD AND WAY OF USE

External.

Preliminary cleaned, the wounded area is covered well with the ointment /by a spatula/ 1-2 times in a day. It is applied until the inflammatory process is calmed, as well as the pain and purification of the wounds.

9. ADVICE FOR PROPER APPLICATION

None.

10. CARENTINE TERM

Meat and organs: null days.

Not allowed for use by animals, whose milk is designed for human consumption.

11. SPECIAL CONDITIONS FOR STORAGE OF THE PTODUCT

To be stored in places, that are out of reach and sight of children.

To be stored below 25 °C.

To be kept out of the light.

To be stored in dry places.

Expiration Term after the first opening of the package: 30 days.

12. SPECIAL WARNINGS

Pregnancy and Lactation:

It may be applied in time of pregnancy.

Not allowed for use in animals, whose milk is designed for human consumption.

Interaction with other veterinary and medical products and other forms of interaction :

Incompatible with iodine, heavy metals and alcaloides.

13. SPECIAL ACTS TO DESTROY UNUSED PRODUCT OR REMAININGS OF IT IF THEY EXIST

Not to be thrown with rubbish or via canal waters.

Ask your veterinary doctor what to do with waste VMP. These acts will help to keep the environment.

14. DATE OF THE LAST REVISION OF THE TEXT

15/01/2015

15. ADDITIONAL INFORMATION

Not all the sizes of the package may be proposed on the market.