

LEFLET FOR USE

Imidocarb VETPROM solution for injection

For cattle, sheep, horses and dogs.

NAME AND COSTANT ADDRESS OF THE LICENCE FOR USE OWNER AND OF THE PRODUSER, IF THEY ARE DIFFERENT

Licence for use Owner and Producer:

VETPROM AG, Bulgaria, town of Radomir, ul. Otec Paisii №26

NAME AND ADDRESS OF THE VETERINARY AND MEDICAL PRODUCT

Imidocarb VETPROM solution for injection / Imidocarb VETPROM solutio pro injectionibus

CONTENT OF THE ACTIVE SUBSTANCE/CES/ AND THE EXCIPIENTS

1 ml drug product contains like active substance 120 mg Imidocarb dipropionate and like excipients Benzyl alcohol and Water for injections.

THERAPEUTIC INDICATIONS

For treatment and profilaxis of babesiosa in cattle, sheep, horses and dogs.

Not to be applied intravenously. Not to be applied in animals with impaired liver or kidney function.

The most common unfavourable reactions include / pain in the time of injecting and weak holinergetic symptoms (salivation,vomitting, nazal leakage). These symptoms disappear in 1-2 hours .

Holinergetic symptoms symptoms may be relieved with the application of atropine sulphate. Other effects,that may be observed are difficult breathing, diarrhea, inflammation in the spot of injecting, lacrimation, sweathing and enxxiety.

KINDS OF ANIMALS TO WHICH THIS VETERINARY AND MEDICAL PRODUCT IS DESIGNED, DOSAGE FOR EVERY KIND OF ANIMAL METHOD AND WAY OF APPLICATION

Dose: Cattle: 1ml/100 kg/m in heterogenous infections, due to anaplasma and babesia-2.5 ml/100 kg m.; Sheep: 0.5 ml/50 kg m.; Horses: 2ml/100kg m (i.m. application is prefered); Dogs: 0.25ml/10 kg.m. In heterogenous infections,due to erlihia and babesia-0.5 ml/10 kg m., 2 doses with interval of 14 days.

Prophylaxis of babesiosa in cattle , horses and dogs: Cattle: 2.5 ml/100 kg m.; Horses: 2 мл/100 kg m.; Dogs: 0.5мл/10 кг.м. Single dose will prevent for more than 4 weeks. In most cases, single dose is enough for complete healing, but in treatment of infections, caused by B. equi in horses are necessary 2 doses with interval of 24 hours.

Way of application:

Subcutaneously or intramuscularly.i.e./s.c.or i.m./

ADVICE FOR PROPER APPLICATION

This veterinary and medical product is designed only for subcutaneous application. Not to be applied venously.

CARENTEN TERM

Cattle: Meat and organs – 90 days. Milk – 5 days.

Sheep: Meat and organs – 90 days.

Not to be used in horses, whose meat is for human consumption.

Not to be used in sheep, whose milk is for human consumption.

SPECIAL CONDITIONS FOR STORAGE OF THE PRODUCT

Not required. It is stored below <25°C, in dry and dark places.

Store in places, out of reach and sight of children!

Not to be used after the expiration date, pointed upon the label!

SPECIAL WARNING(S)

People with proved oversensitivity to imidocarb dipropionat have to avoid the contact with the veterinary and medical product.

SPECIAL ACTS OF DESTROYING THE UNUSED PRODUCT OR REMAININGS OF IT, IF THEY EXIST

Each unused veterinary and medical product or remainings of it have to be destroyed in accordance with the requirements of local legislation.

DATE OF THE LAST REVISION OF THE LEAFLET FOR USE

June, year 2011 .

ADDITIONAL INFORMATION

Veterinary and medical product is proposed in dark glass vials of 50 ml and 100 ml, first hydrolitic class ,according to. Eur.Ph., closed with rubber and aluminium caps.

For each information, conserning this VMP, please contact the local representative of the licence for use.of the owner .

VETPROM AG,

2400 Radomir, Bulgaria, ul. "Otec Paisii" № 26

Tel.: 0777/8-24-93, 8-00-19

FAX: 0777/8-23-91

E-mail: vetprom@abv.bg