

LEAFLET FOR:

JODOPHARM[®] - - foam-forming aerosol for intrauterine application

JODOPHARM[®] - aerosolum spumiferum intrauterinicum

For cows, buffalo caws and mares.

1. NAME AND ADDRESS OF THE AUTHORIZATION HOLDER AND OF THE PRODUCER (IF THEY DIFFERENT)

VETPROM AD, 2400, Radomir, Bulgaria, 26, Otets Paisiy Str.

Tel.: 0777/8-24-93, 8-00-19; Fax. 0777/8-23-91

E-mail: vetprom@abv.bg

2. NAME OF THE VETERINARY MEDICINAL PRODUCT

JODOPHARM[®] - foam-forming aerosol for intrauterine application

JODOPHARM[®] - aerosolum spumiferum intrauterinicum

3. ACTIVE SUBSTANCE AND EXCIPIENTS

Active Substance:

Iodine – 0.18 g/40 g

Potassium iodide – 0.36 g/40 g

Excipients:

Polavax A-31

Dimethyl Sulphoxide

Ethanol (96 per cent)

Propylene Glycol

Forane 134 A

4. THERAPEUTIC INDICATIONS

In chronic first- to third-degree endometritis; in pyometra.

5. CONTRAINDICATIONS

Not to be used in hypersensitive to the active substances animals.

6. ADVERSE REACTIONS

The product causes hyperemia like all other iodine-containing preparations.

If a serious impact or any other effects are registered in the process of using this product.

If a serious impact or other effect, due to the use of this VMP not listed in this leaflet is observed, please inform your veterinarian immediately.

7. ANIMAL SPECIES, FOR WHICH THIS VMP IS DESIGNED

For cows, buffalo cows and mares..

8. RECOMMENDED DOSES FOR EACH OF THE ANIMAL SPECIES, ADMINISTRATION ROUTE

The contents of one monoblock corresponds to one dose. In most cases, a single treatment is sufficient. In more serious cases the treatment can be repeated once or twice in 7-days intervals. In pyometra two monoblocks could be applied as a single dose.

9. ADVICE FOR PROPER ADMINISTRATION

Intrauterine administration after an uterine investigation and a massage via the rectum. Fix the cervix with one hand and the tip – with the other hand. By pressing the button, which is connected to the tube for 10 to 15 seconds the contents of the monoblock is sprayed. In cases of pyometra, the medicinal product is applied after draining the purulent matter from the uterus.

10. WITHDRAWAL PERIOD

Meat and internal organs: zero days.

Milk: zero days.

11. SPECIAL PRECAUTIONS FOR STORAGE OF THE VMP

Keep the medication out of sight and reach of children!

Do not store above 25°C.

Protect from light.

Keep in dry place.

Keep from shock and keep away from heat above 45 °C!

12. SPECIAL PRECAUTIONS FOR USE

Interactions with other veterinary medicinal products and other forms of interactions:

The presence of organic compounds reduces the antibacterial effect.

13. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED VETERINARY MEDICINAL PRODUCT OR WASTE MATERIALS DERIVED FROM THE USE OF SUCH PRODUCTS

The VMP should not be disposed of together with household wastes or sewage. Ask your veterinarian what to do with the no longer necessary VMPs. These measures would help environmental protection.

14. DATE OF THE LAST UPDATE OF THE TEXT

04/03/2013

15. ADDITIONAL INFORMATION

Not all the sizes of the package can be on the market.

Please, contact the local representative of the authorization holder for any information concerning this veterinary medicinal product:

VETPROM AD, 2400, Radomir, Bulgaria, 26, Otets Paisiy Str.

Tel.: 0777/8-24-93, 8-00-19; Fax. 0777/8-23-91

E-mail: vetprom@abv.bg