

## INFORMATION LEAFLET

**RIVANOL 1% ointment**

**RIVANOL 1% unguentum**

Dogs and cats

### 1. NAME AND PERMANENT ADDRESS OF MARKETING AUTHORIZATION HOLDER AND MANUFACTURER, IF DIFFERENT

VETPROM AD,

Bulgaria, Radomir 2400, 26 Otetz Paisiy Street

Tel.: +359777/8-24-93, 8-02-68; Fax +359777/8-23-91

E-mail: [office@vetprom.bg](mailto:office@vetprom.bg)

### 2. NAME OF THE VETERINARY MEDICINAL PRODUCT

**RIVANOL 1% ointment**

**RIVANOL 1% unguentum**

### 3. CONTENTS OF THE ACTIVE SUBSTANCE(S) AND EXCIPIENT(S)

50 g contain

**Active substance:** Ethacridine Lactate 0.5 g

**Excipients:** Zinc Oxide, Wool Fat, White Soft Paraffin и Purified Water

### 4. THERAPEUTIC INDICATIONS

For treatment of wounds, abrasions, open abscesses, infections of the skin and mucous membranes, inflammation of the conjunctiva, the mucous membranes of the genitals

### 5. CONTRAINDICATIONS

Do not use on wounds and skin exposed to sunlight (photosensitivity).

Do not use in case of hypersensitivity to ethacridine.

### 6. UNDESIRABLE ADVERSE REACTIONS

As adverse reactions allergic contact dermatitis and phototoxicity may occur.

The frequency of adverse reactions is determined by the following classification:

- Very common (more than 1 in 10 animals displaying adverse reactions during one course of the treatment)

- Common (more than 1 but less than 10 animals per 100 animals)
- Uncommon (more than 1 but less than 10 animals per 1,000 animals)
- Rare (more than 1 but less than 10 animals per 10,000 animals)
- Very rare (less than 1 animal in 10,000 animals, including casual reports).

If you notice any serious reaction or other effects resulting from the use of the veterinary medicinal product not listed in this leaflet, please immediately inform your veterinarian.

#### **7. ANIMAL SPECIES INDICATED FOR TREATMENT WITH THE VMP**

Dogs and cats.

#### **8. DOSAGE FOR EACH SPECIES, ROUTE AND METHOD OF ADMINISTRATION**

The ointment is applied from one to three times daily. Pre-cleaned, the affected section is smeared by spatula. In case of conjunctivitis it should be administered in a small amount of the outer eye corner, and then the eye to be closed and gently massaged for uniform distribution of the product.

#### **9. ADVICE ON PROPER ADMINISTRATION**

None

#### **10. WITHDRAWAL PERIOD**

Not applicable

#### **11. SPECIAL STORAGE PRECAUTIONS**

Keep out of children

Store at temperature below <25°C.

Keep away from light.

Store in a dry place.

Shelf life after first opening: 30 days.

#### **12. SPECIAL PRECAUTIONS**

None

#### **13. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUTILIZED PRODUCT OR WASTE MATERIALS, IF ANY**

Any unused veterinary medicinal product or waste should be disposed in accordance with local regulations and guidelines. Ask your veterinarian how to dispose VMP. These measures will help to protect the environment.

#### **14. DATE OF LATEST REVISION OF THE TEXT**

26/05/2014

## **15. ADDITIONAL INFORMATION**

Not all package sizes may be found on the market.

For any information about this veterinary product, please contact the local representative of the Marketing Authorisation Holder