

PACKAGE LEAFLET

SULFATHIAZOL 5% ointment 50g

Skin ointment for cattle, horses, sheep, goats, pigs, dogs and cats

1. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER AND OF THE MANUFACTURER IF THEY ARE DIFFERENT

VETPROM AD, Bulgaria,
2400 Radomir, 26 Otets Paisii Str
Tel.: 0777/8-24-93, 8-02-68; fax: 0777/8-23-91
E-mail: vetprom@abv.bg

2. NAME OF THE VETERINARY MEDICINAL PRODUCT

SULFATHIAZOL 5% ointment

3. ACTIVE SUBSTANCE AND EXCIPIENT CONTENT

Content in 50g:

Active substances:

Sulfathiazole Sodium – 2.5g

Excipients:

Cod-liver Oil

Wool Fat

White Soft Paraffin

Purified Water

4. THERAPEUTIC INDICATIONS

For prophylaxis and treatment of clean surgical wounds (cosmetic and prophylactic, orthopedic, abdominal surgeries, castrations), wounds healing by primary intention, and spontaneous wounds – infected or noninfected.

CONTRAINDICATIONS

Do not use in case of hypersensitivity to the active substance or to some of the excipients.

Do not use in the presence of a purulent discharge as its activity would be reduced.

6. ADVERSE REACTIONS

Potential occurrence of skin rashes.

If you notice some serious or other effects of this veterinary medicinal product not listed in this package leaflet, please inform immediately your veterinarian.

7. ANIMAL SPECIES FOR WHICH THIS VETERINARY MEDICINAL PRODUCT IS INTENDED

Cattle, horses, sheep, goats, pigs, dogs and cats

8. DOSAGE FOR EACH ANIMAL SPECIES, METHOD AND ROUTE OF ADMINISTRATION

Dose:

The necrotic and purulent wound areas (affected skin surfaces) should be cleansed . Afterwards the product is applied and carefully rubbed in with gentle movements.

Method of administration:

Topical application to skin.

9. INSTRUCTIONS FOR PROPER ADMINISTRATION

Observe the directions for use.

10. WITHDRAWAL PERIOD

Meat and internal organs: 0(zero) days.

Milk: 0(zero) hours.

11. SPECIAL REQUIREMENTS FOR STORAGE

Keep out of the sight and reach of children.

Store below 25 °C.
Protect from light.
Store in dry places.

Do not use this veterinary medicine after the expiry date which is stated on the label.
Shelf life after first opening: 30 days.

12. SPECIAL WARNINGS

Special warnings for any animal species for which the veterinary medicinal product is intended:

The product is intended for veterinary use only, by a veterinarian.

Pregnancy:

This product can be used during pregnancy.

Lactation:

This product can be used during lactation.

Fertilising ability:

May be used to animals for breeding.

Interactions with other veterinary medicinal products and other forms of interaction

Incompatible with anaesthesin, procaine and dicain .

Overdosage (symptoms, emergency measures, antidotes if required)

In case of overdose, Vitamin B1 and alkaline agents should be used.

Incompatibilities:

Do not mix with anaesthesin, procaine and dicain .

13. SPECIAL PRECAUTIONS FOR DISPOSAL OF ANY UNUSED PRODUCT OR WASTE MATERIAL

Do not throw away any veterinary medicines via wastewater or household waste. Ask your veterinarian how to throw away veterinary medicines you no longer use. These measures will help protect the environment.

14. DATE OF LATEST REVISION OF THE TEXT

15/10/2014

15. FURTHER INFORMATION

For any information in relation to this veterinary medicinal product, please contact the local representative of the marketing authorization holder.

VETPROM AD

2400 Radomir, Bulgaria, 26 Otets Paisii Str

Tel: 0777/8-24-93, 8-00-19

FAX: 0777/8-23-91

E-mail: vetprom@abv.bg