

User guide

TINCTURA IODI
Solutio iodi spirituosa 5%

TINCTURA IODI
Tincture of iodine 5%

For veterinary use only

Name and address of the marketing authorisation holder: VETPROM AD, Radomir, Bulgaria, 26 Otets Paisiy Str.

Tincture iodine 1 ml of the medicinal product contains 0.05 g Iodine and 0.02 g Potassium Iodine as active substances; Ethanol and 96% Purified water as excipients (solvents).

Therapeutic Indications: Intended for disinfection on the surgical field; to mitigate wound (around them), superficial wounds of irregular granulation; with fistulas and ulcers; in actinomycosis, botriomikoza; in trichophytia and other skin mycoses.

Contraindications for use: Do not apply on mucosa; in furunculosis and urticaria.

Adverse reactions: Local irritating. At low concentrations on the skin it causes a burning sensation, prickling, hyperemia with reflex impact on the function of other organs, and in high concentration especially on the mucosa can cause inflammation and necrosis. It penetrates deeply and can provide resorptive effect.

Medicinal or other forms of interactions: Iodine reacts with protein and other organic compounds which reduces its effect. Incompatible with other disinfectants and compounds containing mercury.

Target Species, dose units for each type of animal and the route of administration:

Animals: All animals.

Route of administration: Externally.

The place that is a subject of treatment should be carefully mitigated (mitigate around the wounds). In ulcers and fistulas mitigation is done after pre-cleaning and surgical treatment.

Overdose: There is no risk of overdose due to route of administration.

In case of possible ingestion lacrimation, salivation, nausea, vomiting, diarrhea, catarrh, cough, sometimes lung edema, nephritis and others occurs. Iodine acne and other iodine lesions can be observed.

Management of overdose: Apply starch, flour, sodium thiosulfate, sodium bicarbonate, down from flaxseed, milk, egg whites and symptomatic agents - korazol, kordiamin and others.

Withdrawal period: None.

Special precautions for product storage: Not required.

Storage: At room temperature (15-25°C), dry and protected from light place.

Additional safety data and health protection for the people working with VMP, special precautions and warnings: Not required.

Special precautions for disposal of unused veterinary medicinal product or its wastes: as per stated in the Waste Management Law.

Date of latest revision of the leaflet: January 2008